



Milan S. Chakrabarty, M.D.
General Gastroenterology

Sandra Del Valle, PA-C
Gastroenterology Physician Assistant

Indraneel Chakrabarty, M.D., M.A.
Advanced & Interventional
Gastroenterology

Kathleen Linke, PA-C
Gastroenterology Physician Assistant

Two Locations:

HEMET

1003 E. Florida Avenue · Suite 101 · Hemet, CA 92543 · (951) 652-2252

TEMECULA

31515 Rancho Pueblo Rd. · Suite 202 · Temecula, CA 92592
(951) 383-6001 (By appt.)

GASTROINTESTINAL UPPER GASTROINTESTINAL (GI) ENDOSCOPY (EGD)

What is Upper Gastrointestinal (GI) Endoscopy?

Upper GI endoscopy is a procedure that uses a lighted, flexible endoscope to see inside the upper GI tract. The upper GI tract includes the esophagus, stomach, and duodenum—the first part of the small intestine.

What problems can an Upper GI Endoscopy detect?

Upper GI endoscopy can detect

- ulcers
- precancerous conditions
- inflammation
- abnormal growths
- bowel obstruction
- hiatal hernia

When is an Upper GI Endoscopy used?

Upper GI endoscopy can be used to determine the cause of:

- abdominal pain
- vomiting
- gastric reflux
- anemia
- nausea
- swallowing difficulties
- unexplained weight loss
- bleeding in the upper GI tract

Upper GI endoscopy can be used to remove stuck objects, including food, and to treat conditions such as bleeding ulcers. It can also be used to biopsy tissue in the upper GI tract. During a biopsy, a small piece of tissue is removed for later examination with a microscope.

How to Prepare for an Upper GI Endoscopy

The upper GI tract must be empty before an Upper GI Endoscopy can be performed. Generally, no eating or drinking is allowed for 8 hours before the procedure. Smoking and chewing gum are also prohibited during this time.

Patients should tell their doctor about all health conditions they have—especially heart and lung problems, diabetes,

and allergies—and all medications they are taking. Patients may be asked to temporarily stop taking medications that affect blood clotting or interact with sedatives, which are often given during an Upper GI Endoscopy.


Medications and vitamins that may be restricted before and after upper GI endoscopy include:

- blood thinners
- diabetes medications

Driving is not permitted for 12 to 24 hours after upper GI endoscopy to allow sedatives time to completely wear off. Before the appointment, patients should make plans for a ride home.

How is an Upper GI Endoscopy performed?

Patients may receive a local, liquid anesthetic that is gargled or sprayed on the back of the throat. The anesthetic numbs



the throat and calms the gag reflex. An intravenous (IV) needle is placed in a vein in the arm if a sedative will be given. Sedatives help patients stay relaxed and comfortable. While patients are sedated, the doctor and medical staff monitor vital signs.

During the procedure, patients lie on their back or side on

Continued next page.



an examination table. An endoscope is carefully fed down the esophagus and into the stomach and duodenum. A small camera mounted on the endoscope transmits a video image to a video monitor, allowing close examination of the intestinal lining. Air is pumped through the endoscope to inflate the stomach and duodenum, making them easier to see. Special tools that slide through the endoscope allow the doctor to perform biopsies, stop bleeding, and remove abnormal growths.

Recovery from an Upper GI Endoscopy:

After an Upper GI Endoscopy, patients are moved to a recovery room where they wait about an hour for the sedative to wear off. During this time, patients may feel bloated or nauseated. They may also have a sore throat, which can stay for a day or two. Patients will likely feel tired and should plan to rest for the remainder of the day. Unless otherwise directed, patients may immediately resume their normal diet and medications.

Some results from upper GI endoscopy are available immediately after the procedure. The doctor will often share results with the patient after the sedative has worn off. Biopsy results are usually ready in a few days.

What are the risks associated with an Upper GI Endoscopy?

Risks associated with an Upper GI Endoscopy include:

- abnormal reaction to sedatives
- bleeding from biopsy
- accidental puncture of the upper GI tract that may require surgical correction

Patients who experience any of the following rare symptoms after an Upper GI Endoscopy should contact their doctor immediately:

- swallowing difficulties
- throat, chest, and abdominal pain that worsens
- vomiting
- bloody or very dark stool
- fever

Points to Remember:

- Upper gastrointestinal (GI) endoscopy is a procedure that uses a lighted, flexible endoscope to see inside the upper GI tract.
- To prepare for an Upper GI Endoscopy, no eating or drinking is allowed for 8 hours before the procedure. Smoking and chewing gum are also prohibited.
- Patients should tell their doctor about all health conditions they have and all medications they are taking.
- Driving is not permitted for 12 to 24 hours after an Upper GI Endoscopy to allow the sedative time to wear off. Before the appointment, patients should make plans for a ride home.
- Before an Upper GI Endoscopy, the patient may receive a local anesthetic to numb the throat.
- An intravenous (IV) needle is placed in a vein in the arm if a sedative will be given.
- During an Upper GI Endoscopy, an endoscope is carefully fed into the upper GI tract and images are transmitted to a video monitor.
- A doctor can remove polyps, stop bleeding and biopsy abnormal-looking tissue during an Upper GI Endoscopy by using special tools that slide through the endoscope.
- After an Upper GI Endoscopy, patients may feel bloated or nauseated and may also have a sore throat.
- Unless otherwise directed, patients may immediately resume their normal diet and medications.
- Possible risks of an Upper GI Endoscopy include abnormal reaction to sedatives, bleeding from biopsy, and accidental puncture of the upper GI tract.
- Driving is not permitted for 24 hours after an Upper GI Endoscopy to allow the sedative time to wear off.

Source: National Digestive Diseases Information Clearinghouse (NDDIC) website.